Validity of the Hebrew Version of Profiles of Occupational Engagement in People with Schizophrenia (POES) Assessment Tool

Hadas Hedva Levy, Ariel Zilberstein, Lena Lipskaya-Velikovsky

Hadas Hedva Levy, MSc, OT, Day Care Unit, Psychiatric Division, Rambam Health Care Medical Center. hadasle@gmail.com

Ariel Zilberstein, MD, Psychiatrist, Day Care Unit, Psychiatric Division, Rambam Health Care Medical Center. a_zilberstein@rambam.health.gov.il Lena Lipskaya-Velikovsky, PhD, OT, Occupational Therapy Department, Tel Aviv University. lenasky@gmail.com

Key words: Occupational therapy, mental health, evaluation procedures, patterns of participation

Abstract

Schizophrenia is characterized by psychotic episodes and includes emotional, motivational, cognitive and motor symptoms that influence occupational engagement. Occupational engagement associates with health and well-being, and is considered to be a focus of occupational therapy intervention. The Profiles of Occupational Engagement in people with Schizophrenia (POES) provides a systematic description of occupational engagement and occupational balance, based on a time-use diary completed by people with schizophrenia. The POES has been translated into Hebrew, however, psychometric properties have not been established yet. objectives: To establish criterion and construct validity for the POES Hebrew version. Methods: A cross-sectional comparative study, with 42 participants divided into three groups: (1) people with schizophrenia (N=14), (2) people with personality disorders (N=14) and (3) healthy controls (N=14) matched by age and gender to the first study group. The evaluation procedures included: POES, participation questionnaire, functional capacity and cognitive assessments. Results: Occupational engagement level as evaluated by POES was significantly different between participants with and without psychiatric diagnosis (-4.58<Z<-4.57, p<0.001). Among people with schizophrenia, POES score was associated with cognitive ability (rs=0.566, p<0.05), functional capacity (rs=0.78, p<0.001) and participation diversity (rs=0.733, p<0.01). A further correlation between the first and second measuring of POES was found, while controlling for number of days between the assessments (rpartial=0.577, p<.05). **Discussion and Conclusions:** This study provides initial evidence for discriminant, concurrent and convergent validity of the POES Hebrew version. This supports the use of the POES Hebrew version as a part of the evaluation process by occupational therapists with people with schizophrenia and other psychiatric disorders.